

# Vedafil

## QUICK GUIDE<sup>2</sup>

### WHEN TO TAKE IT

About **1 hour** prior to anticipated sexual activity

---

### WHEN WILL IT START WORKING

Varies from person to person

From **1/2 - 1 hour**

---

### HOW LONG WILL IT WORK FOR

Works only when sexually aroused

Up to **5 hours** when you need it

### FOOD AND ALCOHOL

Drinking alcohol may affect your ability to get an erection, so avoid excessive drinking

**No**, may take longer to work when taken with a meal

---

### MAXIMUM FREQUENCY OF USE

**Once per day** (25 mg or 50 mg or 100 mg tablet)

---

### SIDE EFFECTS<sup>†</sup>

Headache, flushing, nasal congestion, indigestion, diarrhoea, decreased or loss in vision, cardiovascular events

**Vedafil**<sup>®</sup>  
Sildenafil citrate

Scan QR code  
for more information



vedafil.co.nz

## CONNECT REWARDS

Enjoy every

**5<sup>th</sup> PACK** FREE

Terms and conditions to redeem 5<sup>th</sup> pack apply

Scan QR code for more information



[vedafil.co.nz/connect-rewards](https://vedafil.co.nz/connect-rewards)

<sup>1</sup>These are some of the more common or/and very serious side effects. for a comprehensive list consult your healthcare professional or refer to the Consumer Medicine Information for more details.

**References:** 1. IQVIA Data. New Zealand. ATC 3. G04E ERECTILE DYSFUNCTION. Counting Units. MAT January 2024. 2. Vedafil<sup>®</sup> Consumer Medicine Information (20 August 2021). Available at <https://www.medsafe.govt.nz>. Accessed March 2024. **Vedafil<sup>®</sup>** (sildenafil citrate) 25 mg, 50 mg & 100 mg tablets. Prescription Medicine except when supplied by a pharmacist who has successfully completed an approved training programme. For the treatment of adult men with erectile dysfunction (ED). Do not take if you: are allergic to sildenafil or any excipients; taking nitrates or guanylate cyclase stimulators; have risk factors or conditions for which sexual activity is inadvisable; unusually high, low or uncontrolled blood pressure; had a heart attack or stroke in the past 6 months; severe liver condition; certain eye conditions such as NAION or hereditary eye disorders. Possible side effects may include: chest pain, palpitations, fits, prolonged erection, decreased or loss of vision or hearing, allergic reactions. Caution when driving or operating machinery. If symptoms persist or you have side effects, see your healthcare professional. Take strictly as directed. Medicines have benefits and risks. For further information, ask your healthcare professional or view the Consumer Medicine Information at [www.medsafe.govt.nz](http://www.medsafe.govt.nz). Ask your doctor if Vedafil<sup>®</sup> is right for you. Vedafil<sup>®</sup> is an unfunded medicine- doctor's fees and pharmacy charges will apply. Vedafil<sup>®</sup> is a Viatriis company trade mark, Viatriis Limited, Auckland. Copyright© 2024 Viatriis Inc. All rights reserved. VED-2024-0001. TAPS NP20641.

QUICK GUIDE<sup>2</sup>

**Vedafil<sup>®</sup>**  
Sildenafil citrate

**NZ'S #1 SELLING TREATMENT**  
FOR ERECTILE DYSFUNCTION.<sup>1</sup>

  
VIATRIS